

Links of Prime Ideals

Alberto Corso

Claudia Polini*

Wolmer V. Vasconcelos†

Department of Mathematics, Rutgers University
New Brunswick, New Jersey 08903
e-mail: corso, polini, vasconce@math.rutgers.edu

Abstract

We exhibit the elementary but somewhat surprising property that most direct links of prime ideals in Gorenstein rings are equimultiple ideals. It leads to the construction of a bountiful set of Cohen–Macaulay Rees algebras.

1 Introduction

Grosso modo, a link of an ideal $\mathfrak{p}$ of a Noetherian ring R is an ideal of the form $I = (\mathbf{z}): \mathfrak{p}$, where $\mathbf{z} = z_1, \dots, z_g$ is a regular sequence and g is the codimension of $\mathfrak{p}$. This is a very common operation in commutative algebra, particularly in duality theory, and plays an important role in current methods to effect primary decomposition of polynomial ideals (see [2]).

More precisely, let $(R, \mathfrak{m})$ be a Cohen–Macaulay local ring, and let I and J be two ideals. I and J are said to be *linked* if there exists a regular sequence $\mathbf{z} = z_1, \dots, z_g \subset I \cap J$ such that $I = (\mathbf{z}): J$ and $J = (\mathbf{z}): I$. If R is a Gorenstein ring and J is an unmixed ideal, then for any regular sequence $z_1, \dots, z_g \subset J$, where g is the codimension of J , the ideals J and $I = (\mathbf{z}): J$ are linked (see [10]). In this case we simply say that I is a *link* of J .

AMS 1991 *Mathematics Subject Classification*. Primary 13H10; Secondary 13C40, 13D40, 13D45, 13H15.

Key words and phrases. Cohen–Macaulay, Gorenstein ring, canonical module, Koszul homology, linkage of an ideal, equimultiple, syzygetic, Northcott ideal, reduction of an ideal, reduction number, Rees algebra, associated graded ring.

*The author was partially supported by NSF grant STC-91-19999.

†The author was partially supported by NSF grant DMS-93-43436.

To frame our results we recall a notion of Northcott–Rees (see [9]). An ideal $J \subset I$ is a *reduction* of I if $JI^r = I^{r+1}$ for some integer r ; the least such integer is the *reduction number* of I with respect to J . Phrased otherwise, this means that

$$R[Jt] = \sum_{n=0}^{\infty} J^n t^n \xrightarrow{\quad} \sum_{n=0}^{\infty} I^n t^n = R[It]$$

is a finite morphism of the associated Rees algebras, and r is the bound of the degrees required to generate $R[It]$ as a module over $R[Jt]$. If J is generated by a regular sequence the ideal I is said to be *equimultiple*.

There are comparatively few general classes of ideals in Cohen–Macaulay rings whose associated Rees algebra is Cohen–Macaulay. The most basic case, of an ideal I generated by a regular sequence $\mathbf{z} = z_1, \dots, z_g$, has a very explicit presentation as a quotient of a polynomial ring

$$R[It] \simeq R[T_1, \dots, T_g] / I_2 \begin{pmatrix} z_1 & \cdots & z_g \\ T_1 & \cdots & T_g \end{pmatrix},$$

where the relations are the maximal minors of a matrix whose two rows are essentially a set of indeterminates. From [1] it follows that $R[It]$ is Cohen–Macaulay.

Other classes of ideals, with Cohen–Macaulay Rees algebras, include maximal ideals with restrictions on their multiplicities, classical determinantal ideals attached to generic matrices, ideals whose Koszul homology modules have conveniently high depths, and ideals having nice reductions.

The ideals studied here are links of a prime ideal $\mathfrak{p}$. Most of them turn out to be equimultiple, of reduction number one, provided the localization $R_{\mathfrak{p}}$ is a Gorenstein ideal. As a consequence, if R is a Gorenstein ring and $\mathfrak{p}$ is a Cohen–Macaulay ideal then the corresponding associated graded ring will be Cohen–Macaulay. Since ideals such as $\mathfrak{p}$ are very common, the process described here provides Cohen–Macaulay algebras with a great deal of profusion.

2 Equimultiple Ideals

The relationship between a regular sequence and the corresponding link of a prime is discussed next. The general terminology used here is that of [7]; the basic reference for local duality and the theory of the canonical module will be [3].

Two pieces of notation will be used: if R is a local ring and M is a finitely generated R -module, $\nu(M)$ denotes its minimal number of generators; if furthermore M is Artinian, we denote its length by $\lambda(M)$.

Theorem 2.1 *Let R be a Cohen–Macaulay ring, $\mathfrak{p}$ a prime ideal of codimension g , and let $\mathbf{z} = z_1, \dots, z_g \subset \mathfrak{p}$ be a regular sequence. Set $J = (\mathbf{z})$ and $I = J : \mathfrak{p}$. Suppose that $R_{\mathfrak{p}}$ is a Gorenstein ring. Then I is an equimultiple ideal with reduction number one, more precisely,*

$$I^2 = JI$$

in the following two cases:

- (a) $R_{\mathfrak{p}}$ is not a regular local ring;
- (b) $R_{\mathfrak{p}}$ is a regular local ring of dimension at least 2 and two elements in the sequence $\mathbf{z}$ lie in the symbolic square $\mathfrak{p}^{(2)}$.

The proof requires the assemblage of 3 elementary techniques. We deal with them separately first.

Reduction to maximal ideals

To establish the equality $I^2 = JI$, we examine the associated prime ideals of JJ . To that end, looking at the exact sequence (see [5])

$$0 \longrightarrow J/JI \longrightarrow R/JI \longrightarrow R/J \longrightarrow 0,$$

since

$$J/JI = J/J^2 \otimes R/I = (R/I)^g,$$

it follows that the associated primes of JJ satisfy the condition

$$\text{Ass}(R/JI) \subset \text{Ass}(R/I) \cup \text{Ass}(R/J).$$

Therefore all the associated primes of JJ have codimension g , and since I and J are equal at any localization $R_{\mathfrak{q}}$, where $\mathfrak{q}$ is a prime of codimension g , distinct from $\mathfrak{p}$, we may finally assume that R is a local ring and $\mathfrak{p}$ its maximal ideal.

Syzygetic ideals

In making comparisons between the primary ideals I^2 and JJ we consider the following diagram:

$$\begin{array}{ccc}
 & I & \\
 \lambda(I/I^2) \swarrow & & \searrow \lambda(I/J) \\
 I^2 & & J \\
 \lambda(I^2/JI) \swarrow & & \searrow d\lambda(R/I) \\
 & JJ &
 \end{array}$$

We need to establish some relationships between these lengths. This will be done in the next result which contains our main computation.

Proposition 2.2 *Let $(R, \mathfrak{m})$ be a Cohen–Macaulay local ring of Krull dimension d , let $J = (a_1, \dots, a_d)$ be a $\mathfrak{m}$ -primary ideal and let $J \subset I$. Suppose I is minimally generated by $d + r$ elements $(J, b_1, \dots, b_r)$. Denote by H_1 the 1-dimensional Koszul homology module of I and by $\delta(I)$ the kernel of the natural surjection from the symmetric square of I onto I^2 . Then*

$$\lambda(I/J) = \lambda(I^2/JI) + r\lambda(R/I) - \lambda(H_1) + \lambda(\delta(I)). \quad (1)$$

Proof. Let

$$0 \longrightarrow Z \longrightarrow R^{d+r} \longrightarrow I \longrightarrow 0,$$

be a minimal presentation of I . Tensoring with R/I we obtain the exact sequence (see [11])

$$0 \longrightarrow \delta(I) \longrightarrow H_1 \longrightarrow (R/I)^{d+r} \longrightarrow I/I^2 \longrightarrow 0, \quad (2)$$

where $\delta(I)$ accounts for the cycles in Z whose coefficients lie in I .

This leads to $\lambda(I/I^2) - (d + r)\lambda(R/I) + \lambda(H_1) - \lambda(\delta(I)) = 0$, or equivalently

$$\lambda(I/I^2) - d\lambda(R/I) = r\lambda(R/I) - \lambda(H_1) + \lambda(\delta(I)). \quad (3)$$

To account for the left-hand side of (3), consider the following two exact sequences

$$0 \longrightarrow J/JI \longrightarrow I/JI \longrightarrow I/J \longrightarrow 0,$$

$$0 \longrightarrow I^2/JI \longrightarrow I/JI \longrightarrow I/I^2 \longrightarrow 0.$$

From the first one we get that $\lambda(I/JI) = \lambda(J/JI) + \lambda(I/J)$, while from the second one we get that $\lambda(I/JI) = \lambda(I^2/JI) + \lambda(I/I^2)$. By comparing the last two equations, we have that $\lambda(J/JI) + \lambda(I/J) = \lambda(I^2/JI) + \lambda(I/I^2)$, or equivalently

$$\lambda(I/I^2) - \lambda(J/JI) = \lambda(I/J) - \lambda(I^2/JI). \quad (4)$$

As we remarked earlier, J/JI is a free R/I -module of rank d , so from $\lambda(J/JI) = d\lambda(R/I)$, the last equation becomes:

$$\lambda(I/I^2) - d\lambda(R/I) = \lambda(I/J) - \lambda(I^2/JI). \quad (5)$$

If we compare (5) with (3), we get $\lambda(I/J) - \lambda(I^2/JI) = r\lambda(R/I) - \lambda(H_1) + \lambda(\delta(I))$, or equivalently

$$\lambda(I/J) = \lambda(I^2/JI) + r\lambda(R/I) - \lambda(H_1) + \lambda(\delta(I)),$$

as desired. $\square$

Corollary 2.3 *Under the same assumption as in Proposition 2.2, if R is a Gorenstein local ring and $r = 1$ then*

$$\lambda(I/J) = \lambda(I^2/JI) + \lambda(\delta(I)), \quad \text{with } \delta(I) \neq 0. \quad (6)$$

Proof. Since R is Gorenstein and R/I is Cohen-Macaulay, the canonical module of R/I is the last non-vanishing Koszul homology module

$$\omega_{R/I} = \text{Ext}_R^d(R/I, R) = \text{Hom}_{R/J}(R/I, R/J) = (J:I)/J = H_1,$$

as I is generated by a regular sequence plus an extra element. Recall that the canonical module of an Artinian ring is the injective envelope of its residue field and therefore $\lambda(H_1) = \lambda(R/I)$. If we apply Proposition 2.2 we get

$$\lambda(I/J) = \lambda(I^2/JI) + \lambda(\delta(I)).$$

On the other hand, if $\delta(I) = 0$, from (2), we get the short exact sequence

$$0 \longrightarrow H_1 \longrightarrow (R/I)^{d+1} \longrightarrow I/I^2 \longrightarrow 0,$$

which splits since H_1 is injective. Therefore

$$(R/I)^{d+1} \simeq H_1 \oplus I/I^2.$$

Now, $\lambda(H_1) = \lambda(R/I)$ implies that $H_1 \simeq R/I$ and so $I/I^2 \simeq (R/I)^d$, which implies that $\nu(I) = d$ (since $\nu(I) = \nu(I/I^2)$ by Nakayama lemma), contradicting the assumption. $\square$

Remark 2.4 A similar computation would show the following more general formula. Suppose that $(R, \mathfrak{m})$ is a Cohen–Macaulay ring of type $s \geq 2$. Then

$$\lambda(I^2/JI) + \lambda(\delta(I)) = \binom{s+1}{2}.$$

Northcott ideals

Finally, we recall an useful construction introduced by Northcott (see [8]), that provides for a very explicit description of a family of links.

Let R be a Noetherian ring, $\mathbf{u} = u_1, \dots, u_n$ a sequence of elements in R , and $\varphi = (a_{ij})$ a $n \times n$ matrix with entries in R . Denote by $\mathbf{v} = v_1, \dots, v_n$ the sequence

$$[\mathbf{v}]^t = \varphi \cdot [\mathbf{u}]^t.$$

If $\text{grade}(\mathbf{v}) = n$, the ideal $N = (\mathbf{v}, \det \varphi)$ (where $\det \varphi$ is the determinant of the matrix φ) is known as the *Northcott ideal* associated to the sequence $\mathbf{u}$ and the matrix φ . It has then three important properties that can be summarized in the next theorem (see [8]).

Theorem 2.5 *Let $\mathbf{v}$, $\mathbf{u}$ and φ as above. Then either N is the whole ring or it is a proper ideal with the following properties*

- (a) N is a perfect ideal (i.e. its grade is equal to the projective dimension of R/N);
- (b) $(\mathbf{v}):N = (\mathbf{u})$;
- (c) $(\mathbf{v}):(\mathbf{u}) = N$.

The case where $(\mathbf{u})$ is the maximal ideal of a regular local ring is particularly interesting: $\det \varphi$ will then generate the socle of the Artin algebra $R/(\mathbf{v})$.

Proof of Theorem 2.1. According to Corollary 2.3, we must show that $\delta(I) \neq 0$, as this leads to $\lambda(I^2/JI) = 0$. As observed earlier, $\delta(I) = 0$ means that I is also generated by a regular sequence. This puts us in the context of Northcott ideals. Since however $\lambda(I/J) = 1$, it follows that the ideal $N = (J, \det \varphi)$ is perfect and necessarily the maximal ideal of R . In the case that R is not regular, this is a contradiction. In the other case, when at least two of the generators of J lie in the square of the maximal ideal $\mathfrak{m}$, we cannot have the equality $\mathfrak{m} = (J, \det \varphi)$ either. $\square$

3 Cohen–Macaulay Rees Algebras

In this section we are going to use the results obtained in the previous section—in particular Theorem 2.1—in order to study the properties of the Rees algebras of the links of Cohen–Macaulay prime ideals. (We use the terminology and some of the methods described in [4] and some of its references.)

Theorem 3.1 *Let R be a d -dimensional Gorenstein local ring, $\mathfrak{p}$ be a Cohen–Macaulay prime ideal of codimension g , $J = (a_1, \dots, a_g)$ a complete intersection, and let $I = J:\mathfrak{p}$. If one of the following holds:*

- (a) $R_{\mathfrak{p}}$ is not a regular local ring, or
- (b) $R_{\mathfrak{p}}$ is a regular local ring and two of the generators of J belong to $\mathfrak{p}^{(2)}$,

then the associated graded ring $\mathrm{gr}_I(R)$ is Cohen–Macaulay.

Proof. First observe that I is a Cohen–Macaulay ideal (see [10]). Let $R[Jt]$ and $R[It]$ be the Rees algebras of the ideals J and I . Since $I^2 = JI$, by Theorem 2.1, $R[It]$ is a finitely generated $R[Jt]$ -modules and $I \cdot R[Jt] = I \cdot R[It]$.

On the other hand, the exact sequence

$$0 \longrightarrow I \cdot R[Jt] \longrightarrow R[Jt] \longrightarrow R[Jt] \otimes_R R/I \longrightarrow 0$$

implies that $I \cdot R[Jt]$ is a maximal Cohen–Macaulay module over $R[Jt]$, since $R[Jt]$ is a Cohen–Macaulay ring of dimension $d+1$ and the fibre $R[Jt] \otimes_R R/I$ is a polynomial ring in g variables over the Cohen–Macaulay ring R/I ; it is thus a Cohen–Macaulay ring of dimension $\dim(R/I) + g = d - g + g = d$. This means that, when viewed as an $R[It]$ –module, $I \cdot R[It]$ is a maximal Cohen–Macaulay module as well.

The assertion about the associated graded ring of I ,

$$\mathrm{gr}_I(R) = \sum_{n \geq 0} I^n / I^{n+1} = R[It] \otimes_R R/I$$

will follow from depth counting in the exact sequences:

$$0 \longrightarrow I \cdot R[It] \longrightarrow R[It] \longrightarrow \mathrm{gr}_I(R) \longrightarrow 0, \quad (7)$$

$$0 \longrightarrow It \cdot R[It] \longrightarrow R[It] \longrightarrow R \longrightarrow 0, \quad (8)$$

and the canonical isomorphism $It \cdot R[It] \simeq I \cdot R[It]$. $\square$

Corollary 3.2 *If in Theorem 3.1 $d \geq 2$ (which is automatic in the second case), then the Rees algebra of I is Cohen–Macaulay.*

Proof. Since the reduction number of I is 1, the assertion follows from the Theorem and [6]. $\square$

Remark 3.3 Although Theorem 3.1 covers most of the cases of links of prime ideals, the restriction in (b) leaves interesting links out. For example, suppose $\mathfrak{p}$ is a Gorenstein prime ideal of codimension g and let J be a regular sequence in $\mathfrak{p}$ with g elements. The link $I = J : \mathfrak{p}$ is generated by J with an extra element, $I = (J, z)$. If $I_{\mathfrak{p}} = \mathfrak{p}R_{\mathfrak{p}}$, as it may as well occur, then obviously we are not going to have $I^2 = JI$. In fact, in this case, I is going to be generated by analytically independent elements and therefore it will not admit a proper reduction. Nevertheless, one can still show that the Rees algebra of I is Cohen–Macaulay (see [4]).

The authors are also aware of other ideals, in non Gorenstein rings, with some of these properties. But they arise in much more constrained settings.

References

- [1] J. Eagon and D. G. Northcott. Ideals defined by matrices and a certain complex associated with them. *Proc. Royal Soc.* **269** (1962), 188–204.
- [2] D. Eisenbud, C. Huneke and W. V. Vasconcelos. Direct methods for primary decomposition. *Invent. Math.* **110** (1992), 207–235.
- [3] J. Herzog and E. Kunz. *Der kanonische Modul eines Cohen–Macaulay Rings*. Lecture Notes in Mathematics **238** (Springer–Verlag, 1971).
- [4] J. Herzog, A. Simis and W. V. Vasconcelos. Koszul homology and blowing-up rings. In *Commutative Algebra*, Proceedings: Trento 1981 (S. Greco and G. Valla, Eds.), Lecture Notes in Pure and Applied Math. **84** (Marcel Dekker, 1983), 79–169.
- [5] S. Huckaba and C. Huneke. Powers of ideals having small analytic deviation. *Amer. J. Math.* **114** (1992), 367–403.
- [6] S. Ikeda and N. V. Trung. When is the Rees algebra Cohen–Macaulay?. *Comm. Algebra* **17** (1989), 2893–2922.
- [7] H. Matsumura. *Commutative Ring Theory* (Cambridge University Press, 1986).
- [8] D. G. Northcott. A homological investigation of a certain residual ideal. *Math. Ann.* **150** (1963), 99–110.
- [9] D. G. Northcott and D. Rees. Reductions of ideals in local rings. *Proc. Camb. Phil. Soc.* **50** (1954), 145–158.
- [10] C. Peskine and L. Szpiro. Liaison des variétés algébriques. *Invent. Math.* **26** (1974), 271–302.
- [11] A. Simis and W. V. Vasconcelos. The syzygies of the conormal module. *Amer. J. Math.* **103** (1981), 203–224.